



# **Screening of Antimicrobial Activity of Poly Herbal Extracts against Bacterial Pathogens Causing Gastroenteritis in Tharaka Nithi County, Kenya**

**Mbithe P. Kiteme <sup>a\*</sup>, Benson O. Onyango <sup>b</sup>,  
Eric C. Njagi <sup>c</sup> and Fredrick O. Ogolla <sup>a</sup>**

<sup>a</sup> *Department of Biological Sciences, Chuka University, P.O. Box 109-60400, Chuka, Kenya.*

<sup>b</sup> *Department of Biology, Jaramogi Oginga Odinga University of Science and Technology, P.O. Box 210 - 40601 Bondo, Kenya.*

<sup>c</sup> *Department of Physical Sciences, Chuka University, P.O. Box 109-60400, Chuka, Kenya.*

## **Authors' contributions**

*This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.*

## **Article Information**

DOI: 10.9734/SAJRM/2023/v15i2284

## **Open Peer Review History:**

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/98790>

**Original Research Article**

**Received: 15/02/2023**

**Accepted: 18/04/2023**

**Published: 27/04/2023**

## **ABSTRACT**

Herbals have been used for decades to treat gastrointestinal infections across the world, including in Tharaka-Nithi County, Kenya. Crude extracts from herbs have long been prescribed by traditional healers for treatment of typhoid, cholera, and shigellosis. However, the effectiveness of extracts obtained using different extraction solvents such as methanol, acetone, and hexane have not fully been evaluated. This study aimed at evaluating the effect of solvents (Ethanol, hexane, and methanol) on the yield of crude extract from plants (*Erythrina abyssinica*, *Aspilia pluriseta*, *Vangueria infausta*, *Ficus sycomorus*, and *Carissa edulis*), assessing the effect of the solvents on extracted active metabolites, and determining the effect of these solvents on extract's bioactivity

\*Corresponding author: Email: [mbithep@gmail.com](mailto:mbithep@gmail.com);

against *Vibrio cholerae*, *Shigella flexneri*, and *Salmonella typhimurium* and, *Escherichia coli* that served as a model organism for screening plant extracts against Gram-negative bacteria. Crude extracts were obtained by soaking dried ground plant parts in individual solvents, which were then concentrated by a rotary evaporator. The phytochemical screening to detect plant metabolites was done qualitatively. Bioassays to analyze the efficacy of the plant crude extracts against the microbes were carried out in 4×3×3×5 factorial experiment laid out in a completely randomized design. The determination of bioactivity of herbal extracts was carried out using minimal inhibition concentration and minimum bactericidal concentration methods. Data obtained on the bioactivity assay (Count of the bacteria colony forming units) was analyzed using the Kruskal-Wallis test at  $\alpha = 0.05$ , and medians were compared by the Wilcoxon rank sum test in Scientific Analysis System version 9.4. Methanol solvent produced higher quantities of crude extracts for all the herbal samples used with, *Vangueria infausta* producing the highest extract (5.06g). Most phytochemicals were present in Methanolic extracts compared to hexane and acetone extracts. There was a significant ( $p < 0.05$ ) difference in the bioactivity of different herbal plants against bacterial pathogens at different concentrations of crude extract. The efficacy of plant extract increased progressively from 100ppm to 1000ppm concentration. Methanol is recommended for use in the extraction of medicinal plant extracts as it leads to improved potency as compared to hexane and acetone.

**Keywords:** Herbal-plants; antimicrobial-activity; cholera; typhoid; shigellosis; Tharaka-Nithi County.

## 1. INTRODUCTION

Medicine is fundamental to healthcare in management and prevention of illnesses given that healthcare is a basic human need to reduce mortality and disease burdens [1] [2]. One of the key components of the United Nations' Sustainable Development Goals is to ensure universal access to safe, effective, and affordable medication [3]. However, the prevalence of substandard and falsified medicines remains a persistent challenge, compounded by the escalating issue of drug resistance. This presents a critical problem in the field of public health and underscores the urgent need for continued efforts to improve the quality and accessibility of medicines worldwide [4]. Comparative research on alternative antimicrobial agents with greater efficacy is needed to counteract resistant pathogens that are continually selected by current therapeutic regimen [5]. Traditional medication from plant extracts have been in use for years to cure many ailments, thus, herbal are alternative and relatively safer sources of treatment [6]. Traditional medicinal herbs contain bioactive compounds which have few side effects, have low propensity to develop resistance, are less toxic and are associated with improved efficacy [7]. Recent advances in research involving traditional herbal medicine have resulted in identification of several bioactive compounds with antimalarial, anti-cancer and antibiotic properties [8]. Important antimicrobial agents of plant include alkaloids, phenylpropanoids i.e.,

flavonoids and terpenoids that include saponins [9] [10].

Kenya has a wide range of flora with over 7,000 plant species [11]. Up to 70% of the rural populace use home remedies from plant parts as the first source of medicine to treat infections [12] [13]. Different plant parts that include flowers, leaves and fruits are used as sources of home remedies [14]. The current study evaluated the antimicrobial activity of *Carissa edulis* roots, *Aspilia pluriseta* leaves, *Vangueria infausta* leaves and *Erythrina abyssinnica* stem that are traditionally used in treatment of gastrointestinal infections in Tharaka-Nithi County. The health benefits of plants mentioned above are attributed to their richness in phytochemical compounds with physiological effects against gastroenteritis bacteria [15]. However, the phytochemicals present in these crude extracts including those used by herbalists in Tharaka-Nithi County is not fully determined.

Solvents that include water, acetone, hexane, methanol and ethanol, are used to extract bioactive compounds from plant materials [16] [8]. Studies has indicated that solvent used in extraction significantly affects extract's yield, quality, extraction velocity, and bioactivity which is attributed to differences in polarity [16] [17] [18]. Use of ethanol has become popular because beside being safe for infusing food items, it does not corrode container used in addition to being easily recoverable with consistent outcome in bioassays involving crude

extract [19]. Some of the active metabolites that have been extracted using ethanol as solvent are; alkaloids, glycosides, flavonoids, terpenoids, steroids, tannins, saponins, and reducing sugars [20] [19]. Efficiency of alcohol has been attributed to its polarity that resulted in extraction of more active ingredients [21] [20]. As reported by Abarca-Vargas *et al.* [22] and Hikmawanti *et al.* [20], ethanol is considered a protic organic solvent that has a polarity index value of 5.2 and with the dielectric constant of 24.55. Methanol as an extraction solvent has resulted in amino acid, carbohydrate, flavonoids, phenolic compounds, phytosterol, proteins, saponins and tannin [23] [24]. Hexanoic plant crude extracts have been reported not to contain Phenol, Saponins and Carbohydrates [25]. The levels of phenolics and flavonoids is determined by the solvent and has been reported (butanol > methanol > ethyl acetate > hexane > chloroform [24]).

Understanding of the medicinal herbs' aspects such as phytochemical, pharmacological and standardization is necessary for the constitution of dosage in an informed manner [11]. It is therefore necessary to evaluate different solvent concentrations that will produce the most efficacious extract against the test pathogens. In this respect, this study evaluated effect of different organic solvents (Methanol, hexane, and acetone) on the efficacy of crude extract from five plant species (*Carissa edulis* roots, *Aspilia pluriseta* leaves, *Ficus sycomorus* stems, *Erythrina abyssinica* barks and *Vangueria infausta* leaves) against human pathogenic bacteria (*Vibrio cholerae*, *Salmonella typhimurium* and *Shigella flexneri*) associated with gastrointestinal infections. The choice of pathogen species was based on their association with gastroenteritis infection and availability at the Kenya Medical Research Institute (KEMRI), making them the most suitable species for the study.

## 2. MATERIALS AND METHODS

### 2.1 Study Area

The plants used in this study were collected in Tharaka-Nithi County in Kenya which is located at a latitude coordinate of 0°9'25.03''S and longitude coordinates of 37°58'41.48''E. Tharaka-Nithi County has four sub counties; Maara, Tharaka North, Tharaka South and Meru South, covering a total area of 2,662.1 km<sup>2</sup>. The

human population in Tharaka-Nithi County is estimated at 365,330 [26].

### 2.2 Experimental Design

Bioassays to analyze the efficacy of the plant crude extracts against the microbes was carried out in 4x3x3x5 factorial experiment laid out in completely randomized design (CRD) having three factors, with factor A being the microbial pathogens [(At four levels) *Vibrio cholerae*, *Salmonella typhimurium* and *Shigella flexneri* and *E. coli*]. Factor B was the concentrations of crude plant extracts at three levels (1000ppm, 500ppm, 100ppm). Factor C was solvent at three levels (Hexane, acetone and methanol). Lastly factor D was plant extract at 5 levels (*Erythrina abyssinica*, *Aspilia pluriseta*, *Vangueria infausta*, *Ficus sycomorus*, *Carissa edulis*).

### 2.3 Collection of Plant Samples, Extraction of Crude Extracts and Phytochemical Analysis

The commonly used herbs were collected with assistance from traditional herbal dispensers in Tharaka North based on the information gathered through questionnaires on the herbs [27]. Information gathered included the part of the plant used, method of extraction and dose. The plant samples were wrapped in aluminum foils and put in Ziplock bags and transported to Chuka university chemistry laboratories for extraction of the metabolites. In the laboratory, the samples were cleaned using running water and rinsed by use of distilled water. They were then chopped in to small sizes using a sterilized panga (an implement for cutting logs) after which they were left on the bench so as to dry for 21 days. The dried samples were then blended in to soft powder. Extraction of crude extract was performed using methanol, acetone and hexane solvents. 100 grams of each ground sample was weighed and dissolved in 350 ml of methanol, acetone and hexane. They were then allowed to stand for 48 hours so that extraction can take place. Thereafter, filtration was done by use of filter papers number 1. The extracts of the three solvents were then concentrated by use of Rotary evaporator at 40 degrees Celsius and were left to dry in the fume chamber. The concentrated extracts were then weighed using a beam balance and wrapped in aluminum foils, placed in stoppered containers and then transported to KEMRI CDC Kisumu for antimicrobial analysis.

Each plant sample was analyzed for presence of tannins, flavonoids, alkaloids, cardiac glycosides, phenolics, saponins, steroids and terpenoids. Qualitative Test for tannins was done according to Ejikeme *et al.* [28]. Five grams of individual plant powder sample was dissolved in five ml of methanol and then filtered. To the filtrate, 3ml of ferric chloride was added. The occurrence of brownish-green or a blue-black color indicated positive test. Test for flavonoids was done following procedure used by Hossain *et al.* [29] where 2% of the sodium hydroxide was dispensed to each plant sample and followed by addition of dilute hydrochloric acid. The presence of a golden yellow precipitate showed a positive test. Test for saponins was done according to Ejikeme *et al.* [28] where distilled water and each sample filtrate was vigorously shaken till a stable persistent froth was formed. To the mixture, drops of olive oil was added. Formation of an emulsion upon addition of olive oil showed a positive test. Salkowski test as explained by Yadav and Agarwala [30] was used to test for terpenoids presence where three milliliters of each plant extract were dissolved in three milliliters of chloroform. Three milliliters of concentrated sulphuric acid were added where a grayish colour indicated a positive test.

Wagner test procedure used by Ghosh *et al.* [31] was adopted to test for the presence of alkaloids where two milliliters of plant extract was dissolved in two milliliters of ethanol. Two drops of 1M Hydrochloric acid were added to each test tube and boiled for a minute. Three drops of Mayors reagent were added. Formation of a reddish -brown precipitate indicated a positive test. Ferric chloride test as described by Kumar *et al.* [32] was used to test for the presence of phenolics where one milliliter of ferric chloride solution was added to three milliliters of each plant extract sample. Formation of a green colour indicated a positive test for phenolics. Steroids were tested by adding two milliliters of acetic anhydride were added to two milliliters of each plant extract sample followed by addition of two milliliters of sulphuric acid. Formation of a blue or green colour indicated a positive test for steroids. Kellar-Kiliani test as described by Parekh and Chanda 2007 was used to test for glycosides where two milliliters of glacial acetic acid were added to five milliliters of each extract. One milliliter of ferric chloride was then added followed by addition of one milliliter of concentrated sulphuric acid. A green blue coloration of the solution indicated a positive test for glycosides.

## 2.4 Antimicrobial Activity of the Plant Crude Extracts against Test Organisms

### 2.4.1 Preparation of the crude extract, culture media and microbial culture inoculant

Different concentrations of each plant extracts were made dissolving 0.1grams of each crude plant extract in ml 0.2% DMSO making a concentration of 1 g/ml [33]. A 1000 ppm solution was prepared by 1ml of the stock solution into 1000ml of distilled water from which 500ppm and 100ppm concentrations was prepared. The Tryptic Soy Broth (TSB) was prepared using a procedure described in the literature [34] where 15 grams of TSB powder was suspended in 500ml of distilled water and then mixed thoroughly. It was slightly warmed to completely dissolve and then autoclaved at 121°C for 15 minutes prior to cooling at room temperature.

### 2.4.2 Minimum inhibitory concentration and minimum bactericidal concentration of the extracts

Type cultures of the microorganisms used in this study were acquired from KEMRI CDC Kisumu and was sub cultured on Tryptic Soy Broth as per the Clinical Laboratory Standards of US [35]. *Shigella flexneri* and *Vibrio cholerae* which is a member of Enterobacteriaceae family was incubated at a temperature of 35 °C in air for 18 hours while *Salmonella typhimurium* which are anaerobes was incubated at 10% carbon (IV) oxide (CO<sub>2</sub>), 10% hydrogen and 80% nitrogen. Standardization of bacterial cells in the broth was done using 0.5 McFarland standard. The McFarland standard was prepared from 0.5ml of 0.048M BaCl<sub>2</sub> (1.17% w/v) added to 99.5ml of 0.18M sulphuric acid (1% v/v) and stirred [36]. The optical density of the colonies was evaluated using a spectrophotometer and adjusted to equal the 0.5 McFarland standard (10<sup>8</sup> CFU/ml) by adding sterile distilled water [37].

Inoculations was done by adding 20 µL of each bacterial suspensions to individual 100 µL of TSB in a microtiter plate where each of the treatment was replicated three times then incubated for 18 hr. The TSB + DMSO was used while TSB alone as negative control. At the end of the incubation period, A 100 µl of the bacteria under different treatments were cultured by spread plate method on the nutrient agar and was incubated for up to 18 hours. Each bacterium was assayed in different plate

individually. At the end of the re-incubation session, each of the plates was divided into four equal parts using a marker pen and counting of the bacteria CFUs was done for every treatment. The extract concentration at which the CFUs counted was < 10 colonies was assumed to be MBC [38]. The *E. coli*, a common human gastrointestinal tract's bacterium that may also cause gastrointestinal illnesses in humans was used in a screening assay as a model organism for testing the effectiveness of plant extracts against Gram-negative bacteria.

## 2.5 Data Analysis

Data obtained on the bacteria colony forming units (Counts) was analyzed using Kruskal Wallis procedure method  $\alpha = 0.05$  in Statistical Analysis System (SAS) version 9.4. The medians from the results were compared by ranking (Wilcoxon rank sum test) in SAS. Analysis only compared plant extracts and excluded results for positive and negative control set ups.

## 3. RESULTS

### 3.1 Phytochemical Screening of Extracts from Herbal Plants to Treat Gastrointestinal Infections in Tharaka Nithi County

Methanol as solvent produced higher quantities of crude extract for all the herbal samples used. *Vangueria infausta* produced the highest crude extract with methanol (5.06g) while *Aspilia pluriseta* produced lower yield [(0.73g) Table 1].

Several phytochemical compounds were identified from the herbs (Table 2). In methanolic extracts all the seven herbs contained tannins, saponins and phenolic compounds. Flavonoids were only absent in *Ficus sycomorus*, terpenoids were absent only in *Aspilia pluriseta* and *Vangueria infausta* leaves. Alkaloids were present in *Carissa edulis*, *Vangueria infausta* leaves and barks. Steroids were present in *Ficus sycomorus*, *Carissa edulis*, *Erythrina abyssinica* barks only while *Vangueria infausta* leaves were the only containing glycosides.

### 3.2 Effect of Solvents Choice and Plant on Efficacy of Plant's Bacterial Colony Forming Units

The selection of solvent had a significant impact ( $p < 0.05$ ) on the colony forming units (cfu) of various bacterial species (*E. coli*, *Vibrio cholerae*,

and *Shigella flexneri*) when treated with different plant extracts. *Erythrina abyssinica* extract resulted in reduced cfu in *E. coli* (Median = 0) and *V. cholerae* (Median = 6) when treated with methanoic and acetonioic extract, respectively. *Aspilia pluriseta* extract resulted in lower cfu in *E. coli* (Median = 3) and *S. flexineri* (Median = 28) when treated with methanoic extract only. *Vangueria infausta* extract resulted in lower cfu in *Salmonella typhimurium* (Median = 17) when treated with methanoic extract. *Ficus sycomorus* extract resulted in lower cfu in *V. cholerae* (Median = 30) and *S. flexineri* (Median = 27) when treated with methanoic extract. Finally, *Carissa edulis* extract resulted in lower cfu in *S. flexineri* (Median = 30) when treated with hexanoic extract only (Table 3).

There was a significant ( $p < 0.05$ ) effect of exposing test bacteria to various plant extracts at different concentrations on the bacteria colony forming units formed (Table 4). Exposing *E. coli* to 1000ppm plant extracts resulted in significantly reduced bacteria cfu counts in methanoic extracts of *Erythrina abyssinica* (Median = 0 cfu), *Aspilia pluriseta* (Median = 3 cfu), and *Ficus sycomorus* (Median = 0 cfu), while remaining significantly higher in hexanoic extract of *Aspilia pluriseta* (Median = 276 cfu) and *Vangueria infausta* (Median = 250 cfu). Exposing *Salmonella typhimurium* to 1000ppm of plant extracts resulted in significantly lower bacteria cfu counts in methanolic extracts of *Ficus sycomorus* (Median = 13 cfu) and *Vangueria infausta* (Median = 17 cfu), while remaining significantly higher in hexanoic (Median = 256 cfu) and acetonioic (Median = 256 cfu) extracts of *Vangueria infausta*, respectively. Exposing *V. cholerae* to 1000ppm of plant extracts resulted in significantly lower bacteria cfu counts in acetonioic extracts of *Erythrina abyssinica* (Median = 0 cfu) and *Aspilia pluriseta* (Median = 1 cfu), while significantly higher was observed in hexanoic (Median = 265 cfu) extract of *Vangueria infausta*. Exposing *Shigella flexneri* to 1000ppm of plant extracts resulted in significantly lower bacteria cfu counts in acetonioic extracts of *Erythrina abyssinica* (Median = 2 cfu) and methanolic extract of *Ficus sycomorus* (Median = 1 cfu), and significantly higher cfu was observed in hexanoic (Median = 278 cfu) extract of *Vangueria infausta*.

Exposing *E. coli* to 500ppm plant extracts resulted in significantly reduced bacteria cfu counts in methanolic extracts of *Erythrina abyssinica* (Median = 0 cfu) and *Aspilia pluriseta* (Median = 3 cfu) but remained significantly higher in hexanoic extract of *Vangueria infausta*

(Median = 258 cfu). Exposure of *Salmonella typhimurium* to 500ppm of plant extracts resulted in significantly reduced bacteria cfu counts in methanolic extracts of *Vangueria infausta* (Median = 62 cfu) but remaining significantly higher in hexanoic (Median = 266 cfu) extracts of *Vangueria infausta*. Exposing *V. cholerae* to 500ppm of plant extracts resulted in significantly reduced cfu counts in acetonic extracts of *Ficus sycomorus* (Median = 4 cfu) and *Aspilia pluriseta* (Median = 5 cfu) while bacteria cfu counts were significantly higher in hexanoic (Median = 287 cfu) extract of *Vangueria infausta*. Exposing *Shigella flexneri* to 500ppm of plant extracts resulted in reduced bacteria cfu counts in methanolic extracts of *Carissa edulis* (Median = 27 cfu) and higher cfu was observed in acetonic (Median = 265 cfu) extract of *Aspilia pluriseta* (Table 4).

When *E. coli* was exposed to 100ppm plant extracts, significantly reduced bacteria cfu counts

occurred in methanolic extracts of *Aspilia pluriseta* (Median = 0 cfu) but remained significantly higher in hexanoic extract of *Vangueria infausta* (Median = 296 cfu). Exposure of *Salmonella typhimurium* to 100ppm of plant extracts resulted in significantly reduced cfu counts in acetonic extracts of *Erythrina abyssinica* (Median = 22 cfu) but remaining significantly higher (Median = 284 cfu) in hexanoic extracts of *Vangueria infausta*. Exposing *V. cholerae* to 100ppm of plant extracts resulted in significantly reduced cfu counts in acetonic extracts of *Aspilia pluriseta* (Median = 9 cfu) while significantly higher in hexanoic (Median = 265 cfu) extract of *Vangueria infausta*. Exposing *Shigella flexneri* to 100ppm of plant extracts resulted in reduced cfu counts in acetonic extracts of *Vangueria infausta* (Median = 22 cfu) and higher bacteria cfu was observed in hexanoic (Median = 288 cfu) extract of *Vangueria infausta* (Table 4).

**Table 1. Mass of crude extracts obtained using methanol, acetone and hexane solvents**

Plant sample	Hexane	Acetone	Methanol
<i>Carissa edulis</i>	1.81g	3.7g	4.02g
<i>Aspilia pluriseta</i>	0.45g	0.6g	0.73g
<i>Physalis peruviana</i>	1.25g	2.4g	3.74g
<i>Vangueria infausta</i>	1.32g	1.13g	5.06g
<i>Erythrina abyssinnicca</i>	0.06g	1.61g	1.76g
<i>Ficus sycomorus</i>	0.5g	1.22g	1.6g

**Table 2. Phytochemicals present in crude extracts of some medicinal plants used for treatment of gastrointestinal infections**

Sample	Tnns <sup>a</sup>	Flv <sup>b</sup>	Spnn <sup>c</sup>	Tpn <sup>d</sup>	Alk <sup>e</sup>	Phen <sup>f</sup>	Ster <sup>g</sup>	Glyc <sup>h</sup>
<b>Methanolic extracts</b>								
<i>Fs</i> <sup>1</sup>	+	-	+	+	-	+	+	-
<i>Ce</i> <sup>2</sup>	+	+	+	+	+	+	+	-
<i>Ap</i> <sup>3</sup>	+	-	+	-	-	+	-	-
<i>Ea</i> <sup>4</sup>	+	+	+	+	-	+	+	-
<i>Vil</i> <sup>5</sup>	+	+	+	-	+	+	-	+
<b>Hexanoic extracts</b>								
<i>Fs</i> <sup>1</sup>	+	-	+	-	-	+	-	-
<i>Ce</i> <sup>2</sup>	+	+	+	+	-	-	+	-
<i>Ap</i> <sup>3</sup>	+	-	-	-	-	+	-	-
<i>Ea</i> <sup>4</sup>	+	+	+	-	-	+	+	-
<i>Vil</i> <sup>5</sup>	+	+	+	-	-	-	-	+
<b>Acetonic extracts</b>								
<i>Fs</i> <sup>1</sup>	+	-	+	-	-	-	-	-
<i>Ce</i> <sup>2</sup>	+	+	+	-	-	-	+	-
<i>Ap</i> <sup>3</sup>	+	-	-	-	-	-	-	-
<i>Ea</i> <sup>4</sup>	+	+	+	-	-	+	-	-
<i>Vil</i> <sup>5</sup>	+	+	-	-	+	-	-	+

<sup>1</sup>*F. sycomorus*, <sup>2</sup>*C. edulis*, <sup>3</sup>*A. pluriseta*, <sup>4</sup>*E. abyssinnica* (barks). <sup>5</sup>*V. infausta* (leaves). <sup>a</sup>Tannins, <sup>b</sup>Flavonoids, <sup>c</sup>Saponins, <sup>d</sup>Terpenoids, <sup>e</sup>Alkaloids, <sup>f</sup>Phenolics, <sup>g</sup>Steroids, <sup>h</sup>Glycosides

**Table 3. Effect of different solvents on efficacy plant's extracts against bacteria growth**

Plant extract	Test bacteria	Hex <sup>6</sup>	Ace <sup>7</sup>	Met <sup>8</sup>	Kruskal-Wallis statistics			
					N	H	df	P value
Eryth <sup>1</sup>	E <sup>k</sup>	217 <sup>a</sup>	200 <sup>a</sup>	0 <sup>b</sup>	9	19.376	2	<.0001
	S <sup>t</sup>	187	177	167	9	2.084	2	0.352
	V	211 <sup>a</sup>	6 <sup>c</sup>	154 <sup>b</sup>	9	19.323	2	<.0001
	S <sup>f</sup>	199 <sup>a</sup>	56 <sup>b</sup>	146 <sup>a</sup>	9	19.142	2	<.0001
Aspi <sup>2</sup>	E <sup>k</sup>	276 <sup>a</sup>	232 <sup>a</sup>	3 <sup>b</sup>	9	17.801	2	0.0001
	S <sup>t</sup>	178 <sup>a</sup>	165 <sup>a</sup>	146 <sup>b</sup>	9	15.737	2	0.0004
	V	200 <sup>a</sup>	1 <sup>c</sup>	7 <sup>b</sup>	9	20.544	2	<.0001
	S <sup>f</sup>	222 <sup>a</sup>	166 <sup>b</sup>	28 <sup>b</sup>	9	13.302	2	0.0013
Vang <sup>3</sup>	E <sup>k</sup>	250 <sup>c</sup>	243 <sup>b</sup>	187 <sup>a</sup>	9	19.441	2	<.0001
	S <sup>t</sup>	256 <sup>a</sup>	234 <sup>b</sup>	17 <sup>c</sup>	9	20.538	2	<.0001
	V	265 <sup>a</sup>	213 <sup>b</sup>	187 <sup>b</sup>	9	19.413	2	<.0001
	S <sup>f</sup>	278 <sup>a</sup>	220 <sup>b</sup>	187 <sup>b</sup>	9	20.711	2	<.0001
Ficus <sup>4</sup>	E <sup>k</sup>	55 <sup>a</sup>	10 <sup>b</sup>	10 <sup>b</sup>	9	12.622	2	0.0018
	S <sup>t</sup>	140	123	111	9	6.091	2	0.05
	V	87	109	30	9	2.309	2	0.3153
	S <sup>f</sup>	79 <sup>b</sup>	111 <sup>a</sup>	27 <sup>b</sup>	9	12.169	2	0.0023
Cari <sup>5</sup>	E <sup>k</sup>	65 <sup>c</sup>	155 <sup>a</sup>	143 <sup>b</sup>	9	16.414	2	0.0002
	S <sup>t</sup>	198 <sup>a</sup>	168 <sup>b</sup>	148 <sup>c</sup>	9	18.618	2	<.0001
	V	30 <sup>b</sup>	176 <sup>a</sup>	148 <sup>a</sup>	9	6.909	2	0.0316
	S <sup>f</sup>	198 <sup>a</sup>	156 <sup>b</sup>	140 <sup>b</sup>	9	12.384	2	0.002

<sup>a</sup>Median values followed by the same letters in rows are not significantly different at  $\alpha = 0.05$ . <sup>1</sup> *Erythrina abyssinica*, <sup>2</sup> *Aspilia pluriseta*, <sup>3</sup> *Vangueria infausta*, <sup>4</sup> *Ficus sycomorus*, <sup>5</sup> *Carissa edulis*, <sup>k</sup> *E. coli*, <sup>t</sup> *S. typhimurium*, <sup>v</sup> *V. cholerae*, <sup>f</sup> *S. flexineri*, <sup>6</sup> Hexane extract, <sup>7</sup> Acetonic extract, <sup>8</sup> Methanoic extra

#### 4. DISCUSSION AND CONCLUSION

##### 4.1 Phytochemical Extraction Using Different Solvent and Analysis of Crude Extracts

Higher yield of phytochemicals were obtained when methanol was used as solvent. These observations are comparable to other results obtained in related studies such as those of Do et al (2014) on the extraction yield of *Limophila aromatica* and that of Kuppusamy et al. [39] on *Commelina nudiflora*. Further, these results are supported by studies which have demonstrated that the efficiency of extraction process is dependent on type of the solvent used [40] [41]. Variation on the amount of extract obtained by different solvent is attributed to difference in their polarity [42]. This could be that the medicinal herbs contain a higher number of polar compounds. High polar compounds are soluble in solvents that have high polarity like in the case of methanol [42]. Liu et al. [43] describes methanol, a polar solvent as a solvent with improved efficiency of solvation as a result of interaction of hydrogen bonds of the polar sites in the metabolites and the solvent than the non-polar.

The phytochemical analysis results showed the occurrence of secondary metabolites such as phenols, saponins, tannins, flavonoids, terpenoids and glycosides which are known for medicinal attributes as well as physiological activities of the herb [44]. These phytochemicals may be produced in response to attack or infection of the plant by the microbes. The phytochemicals have been observed to have antimicrobial properties against numerous micro-organisms [45] [46]. Extracts from *Carissa edulis* had compounds such as terpenoids, tannins, steroids flavonoids, saponins, alkaloids, phenolics and absence of glycosides. The current results are in line with other related studies such as those of Nawaz [47] who reported occurrence of such compounds in stem bark of *Ficus sycomorus*. The results are further in agreement with a study carried out by Madivoli et al. [48].

Extracts from *Ficus sycomorus* had tannins, saponins, terpenoids, phenolics, steroids while flavonoids, alkaloids and glycosides were absent. The occurrence of these metabolites was similar to those documented by Nawaz et al. [47] who observed tannins, flavonoids, saponins and phenolics. This was also in agreement with a

study carried out in Sudan by Osama and Awdelkarim [49]. However, in their study flavonoids and alkaloids were present, this difference could be linked to associated to the difference in climatic conditions or even the soil type in which this herb grow from where the samples were collected.

*Aspilia pluriseta* extracts had tannins, saponins and phenolics while flavonoids, terpenoids, alkaloids, steroids and glycosides were absent. A comparable investigation was conducted by Njeru and Muema [50] within the Mbeere community, Embu County, exploring its potential therapeutic use for tuberculosis. The results of this study differed with those carried out around Maseno University, Kenya by Musyimi et al. [51] who found the presence of alkaloids, steroids and glycosides which were absent in this study. In both studies methanol as a solvent was used for extraction, therefore the difference could be attributed to different rainfall amount or even the soil type in these different regions. *Erythrina abyssinnica* extracts had tannins, flavonoids, saponins, terpenoids, phenolics, steroids while alkaloids and glycosides were absent. These results are supported by a similar study conducted in Tharaka-Nithi county on utilization of herbal medicine among children under five by Nzuki [52] who identified presence of tannins, terpenoids, alkaloids, phenolics and flavonoids.

*Vangueria infausta* leaf extract had tannins, flavonoids, saponins, alkaloids, phenolics and glycosides while terpenoids and steroids were absent. These results agree with report of Obakiro et al. [53] who observed the presence of flavonoids, alkaloids, phenolics and saponins. These results also are in line with results from a study carried in Zimbabwe by Manyarara et al. [54] where the extract is used in treatment of candidiasis. Saponins are present in almost all medicinal herbs which grounds for their curative potential. Biological role of phytochemicals in plants is solely to protect them against invasion by injurious pathogens and organisms. That notwithstanding, certain saponins have been reported to be toxic to endotherms but such toxicity is lower towards mammals. Since they are toxic to quite a number of organisms, saponins find their use in pharmacological, antibiotic and fungicidal applications [55]. Flavonoids which are based on a 15-carbon skeleton is comprised of two benzene ring connected through a heterocyclic pyrane ring. The basic flavonoid structure is the aglycone. These flavonoids have many biochemical properties including antioxidant property which is

governed by function group arrangement. The substitution, configuration and the total number of hydroxyl groups determines the antioxidant property mechanisms such as the ability to chelate metal ions and in scavenging of radical [56] [57]. The mechanism of antioxidant action involves upregulation or protection of antioxidant defences, suppression of injurious reactive oxygen species which is formed by either inhibition of the enzymes or through chelation of the trace elements that are involved in free radical generation and also by scavenging reactive oxygen species [46].

Studies by Meng et al. [58] as well as that by Kumar and Pandey [59] show that flavonoids are present in herbal medicines and are reported to be effective as cardio protective agents, antibacterial, antioxidants and an important part of pharmaceutical and medical applications. In plants, the living and non-living factors bring about production of reactive oxygen species (ROS) leading to oxidative stress [59]. The production of flavonoids is improved as a result of this oxidative stress and they exhibit the ability to hinder ROS generation, ability to quench ROS immediately they are formed [59]. Further, they have the potential of absorbing the energetic solar wavelengths. The antioxidant capability as well as the ability of absorbing ultra violet wavelengths depend on how different rings of flavonoids are substituted [59]. According to Cazorolli et al. [60], flavonoids inhibit enzymes-based activities including peroxidase and those of xanthine oxidase which take part in generation of free radical thus reducing the oxidative damage brought about by the macromolecules.

The phenolic compounds are among the highest and common plant phytochemicals [61]. Phenolic compounds have anti-inflammatory inflammation properties, anti-aging properties, and have the advantage of improvement of endothelial function and causing the inhibition of the cell proliferation and angiogenesis activities [62]. Phenolic compounds accounts for the antioxidant properties exhibited by most plants used as herbal medicine [63]. Phenolic compounds in plants comprises of chemicals like flavonoids, tocopherols and phenolic acids [63] and are available in almost all plant parts. Phenolics have roles such as scavenging of free radicals, have anticarcinogenic effect as well as antioxidant properties. Phenolics are useful in prevention of human chronic diseases like those associated with bacterial as well as parasitic infections [64].



**Table 4. Effect of different solvents and extract concentrations on efficacy of plant's extracts against bacteria growth**

Plant extract	Sol	N	Plant extracts at 1000ppm					Kruskal-Wallis statistics		
			Eryth <sup>1</sup>	Aspi <sup>2</sup>	Vang <sup>3</sup>	Fic <sup>4</sup>	Cari <sup>5</sup>	H	Df	p-value
E <sup>k</sup>	Hex <sup>6</sup>	3	217 <sup>b</sup>	276 <sup>a</sup>	250 <sup>a</sup>	57 <sup>c</sup>	21 <sup>c</sup>	12.433	4	0.014
	Ace <sup>7</sup>	3	154 <sup>b</sup>	232 <sup>a</sup>	243 <sup>a</sup>	11 <sup>c</sup>	165 <sup>b</sup>	11.750	4	0.019
	Met <sup>8</sup>	3	0 <sup>c</sup>	3 <sup>c</sup>	187 <sup>a</sup>	3 <sup>c</sup>	80 <sup>b</sup>	11.835	4	0.018
S <sup>t</sup>	Hex <sup>6</sup>	3	163 <sup>b</sup>	178 <sup>b</sup>	256 <sup>a</sup>	67 <sup>c</sup>	189 <sup>b</sup>	11.145	4	0.025
	Ace <sup>7</sup>	3	22 <sup>d</sup>	165 <sup>b</sup>	234 <sup>a</sup>	122 <sup>c</sup>	168 <sup>b</sup>	12.833	4	0.012
	Met <sup>8</sup>	3	167 <sup>a</sup>	146 <sup>a</sup>	17 <sup>c</sup>	13 <sup>b</sup>	145 <sup>a</sup>	11.212	4	0.024
V <sup>l</sup>	Hex <sup>6</sup>	3	144 <sup>c</sup>	200 <sup>b</sup>	265 <sup>a</sup>	57 <sup>d</sup>	8	13.524	4	0.009
	Ace <sup>7</sup>	3	0 <sup>d</sup>	1 <sup>d</sup>	213 <sup>a</sup>	122 <sup>c</sup>	176 <sup>b</sup>	13.099	4	0.011
	Met <sup>8</sup>	3	164 <sup>b</sup>	7 <sup>c</sup>	220 <sup>a</sup>	7 <sup>c</sup>	38 <sup>c</sup>	12.077	4	0.016
S <sup>f</sup>	Hex <sup>6</sup>	3	144 <sup>c</sup>	222 <sup>b</sup>	278 <sup>a</sup>	28 <sup>d</sup>	200 <sup>b</sup>	13.257	4	0.010
	Ace <sup>7</sup>	3	2 <sup>d</sup>	166 <sup>b</sup>	202 <sup>a</sup>	110 <sup>c</sup>	20 <sup>d</sup>	12.9	4	0.011
	Met <sup>8</sup>	3	109 <sup>b</sup>	28 <sup>d</sup>	187 <sup>a</sup>	4 <sup>d</sup>	89 <sup>c</sup>	12.923	4	0.012
<b>Plant extracts at 500ppm</b>										
E <sup>k</sup>	Hex <sup>6</sup>	3	215 <sup>b</sup>	234 <sup>a</sup>	258 <sup>a</sup>	47 <sup>c</sup>	65 <sup>c</sup>	11.479	4	0.022
	Ace <sup>7</sup>	3	200 <sup>a</sup>	188 <sup>a</sup>	222 <sup>a</sup>	11 <sup>d</sup>	167 <sup>c</sup>	10.444	4	0.034
	Met <sup>8</sup>	3	0 <sup>c</sup>	0 <sup>c</sup>	198 <sup>a</sup>	10 <sup>c</sup>	143 <sup>b</sup>	13.68	4	0.008
S <sup>t</sup>	Hex <sup>6</sup>	3	187 <sup>b</sup>	233 <sup>a</sup>	266 <sup>a</sup>	140 <sup>c</sup>	203 <sup>b</sup>	11.979	4	0.017
	Ace <sup>7</sup>	3	177 <sup>a</sup>	200 <sup>a</sup>	195 <sup>a</sup>	133 <sup>b</sup>	177 <sup>a</sup>	10.627	4	0.031
	Met <sup>8</sup>	3	133 <sup>a</sup>	117 <sup>a</sup>	62 <sup>c</sup>	111 <sup>b</sup>	136 <sup>a</sup>	10.961	4	0.027
V <sup>l</sup>	Hex <sup>6</sup>	3	234 <sup>b</sup>	225 <sup>b</sup>	287 <sup>a</sup>	89 <sup>c</sup>	30 <sup>d</sup>	11.979	4	0.017
	Ace <sup>7</sup>	3	6 <sup>b</sup>	5 <sup>b</sup>	189 <sup>a</sup>	4 <sup>b</sup>	176 <sup>a</sup>	10.627	4	0.031
	Met <sup>8</sup>	3	123 <sup>b</sup>	132 <sup>a</sup>	152 <sup>a</sup>	89 <sup>c</sup>	148 <sup>a</sup>	10.961	4	0.027
S <sup>f</sup>	Hex <sup>6</sup>	3	238 <sup>b</sup>	223 <sup>b</sup>	278 <sup>a</sup>	80 <sup>d</sup>	185 <sup>c</sup>	10.252	4	0.036
	Ace <sup>7</sup>	3	41 <sup>c</sup>	265 <sup>a</sup>	201 <sup>b</sup>	132 <sup>c</sup>	172 <sup>b</sup>	12.533	4	0.013
	Met <sup>8</sup>	3	146	154	142	27	130	8	4	0.092
<b>Plant extracts at 100ppm</b>										
E <sup>k</sup>	Hex <sup>6</sup>	3	258 <sup>a</sup>	276 <sup>a</sup>	296 <sup>a</sup>	64 <sup>b</sup>	74 <sup>b</sup>	11.487	4	0.022
	Ace <sup>7</sup>	3	217 <sup>b</sup>	283 <sup>a</sup>	208 <sup>b</sup>	6 <sup>d</sup>	99 <sup>c</sup>	12.833	4	0.012
	Met <sup>8</sup>	3	3 <sup>d</sup>	0 <sup>d</sup>	87 <sup>b</sup>	38 <sup>c</sup>	159 <sup>a</sup>	13.597	4	0.008
S <sup>t</sup>	Hex <sup>6</sup>	3	220 <sup>b</sup>	266 <sup>a</sup>	284 <sup>a</sup>	160 <sup>c</sup>	189 <sup>b</sup>	12.022	4	0.017
	Ace <sup>7</sup>	3	22 <sup>d</sup>	165 <sup>b</sup>	234 <sup>a</sup>	122 <sup>c</sup>	168 <sup>b</sup>	12.833	4	0.012
	Met <sup>8</sup>	3	204 <sup>b</sup>	267 <sup>a</sup>	202 <sup>b</sup>	115 <sup>d</sup>	166 <sup>c</sup>	12.433	4	0.014
V <sup>l</sup>	Hex <sup>6</sup>	3	211 <sup>b</sup>	265 <sup>a</sup>	89 <sup>c</sup>	188 <sup>b</sup>	8 <sup>d</sup>	13.233	4	0.010
	Ace <sup>7</sup>	3	76 <sup>d</sup>	9 <sup>e</sup>	243 <sup>a</sup>	113 <sup>c</sup>	178 <sup>b</sup>	13.548	4	0.008
	Met <sup>8</sup>	3	193	154	162	27	171	8.956	4	0.062
S <sup>f</sup>	Hex <sup>6</sup>	3	267 <sup>a</sup>	279 <sup>a</sup>	288 <sup>a</sup>	103 <sup>c</sup>	198 <sup>b</sup>	11.215	4	0.024
	Ace <sup>7</sup>	3	84 <sup>c</sup>	76 <sup>c</sup>	22 <sup>d</sup>	110 <sup>b</sup>	156 <sup>a</sup>	12.9	4	0.012
	Met <sup>8</sup>	3	189 <sup>a</sup>	167 <sup>a</sup>	181 <sup>a</sup>	120 <sup>b</sup>	165 <sup>a</sup>	12.714	4	0.013

<sup>a</sup>Median values followed by the same letters in rows are not significantly different at  $\alpha = 0.05$ . <sup>1</sup> *Erythrina abyssinica*, <sup>2</sup> *Aspilia pluriseta*, <sup>3</sup> *Vangueria infausta*, <sup>4</sup> *Ficus sycomonus*, <sup>5</sup> *Carissa edulis*, <sup>k</sup> *E. coli*, <sup>t</sup> *S. typhimurium*, <sup>l</sup> *V. cholerae*, <sup>f</sup> *S. flexineri*, <sup>6</sup> Hexane extract, <sup>7</sup> Acetonic extract, <sup>8</sup> Methanoic extract

Steroids have been reported to be endowed with antibacterial phytochemicals [65]. Further, the steroids are considered to be essential phytochemicals as they are associated with sex hormones [30]. Despite having cytotoxic properties, steroids have medical significance [30]. Reports indicate that alkaloids obtained from plants are used as therapy for anaesthetics, antioxidants, antidiabetic, antimicrobial and muscle relaxants. In plants, alkaloids offer

protection against injurious singlet reactive oxygen which may be produced in plant tissues and excited by light source [66]. According to report by Ayoola *et al.* [67], alkaloids possess anti-bacterial properties that are exploitable for treatment of bacterial related diseases. Additionally, alkaloids have Anti spasmodic as well as the analgesic significance [67]. Tannins comprises of complex derivatives of the acid (polyhydroxy benzoic) an organic or non-

nitrogenous compound endowed with the ability of precipitating proteins. They are capable of dissolving in water to form colloidal solutions. Further, tannins are soluble in acetone alcohol but are however slightly soluble in organic solvents including chloroform and ethyl acetate. Presence of groups such as carboxylic as well as the free phenolic gives tannins solutions acidic characteristics [68]. Tannins are medicinally used as antidiarrheal, as haemostatic and as antihemorrhoidal metabolites. Due to their anti-inflammatory activities, are useful in the treatment of enteritis, bowel disorders and gastritis. Further, tannins may be used to inhibit viral infections [69], bacterial infection as well as to treat parasitic infections [70]. According to Awuchi [71] glycosides are molecules that have their sugar attached to a different functional group by way of glycosidic bond. Reports by Njoku and Obi [72] indicate that glycosides lower blood pressure.

#### 4.2 Effect of Solvents Choice and Plant on Efficacy of Plant's Bacterial Colony Forming Units

The selection of solvent used in extraction of phytochemicals had a significant impact on the colony forming units (CFUs) of various bacterial species. According to Korir *et al.* [73]. *Erythrina abyssinica* extracts are active against gram positive bacteria as well as on the gram-negative bacteria but have lower efficacy against *E. coli*. Differences in response of different bacteria species towards extracts of some plant species have been reported in other work [74]. Specifically, evaluation of antimicrobial activity of *Ficus* sp extracts inhibit different bacteria species at different levels [75]. Variability of plant extracts in reducing the growth of bacterial colony units in this study could be attributed to plants species with different strength of phytochemicals. This argument is supported by related work of Odunbaku *et al.* [76] and Chilufya *et al.* [77] on the minimal inhibition of sycomorous. According to Mostafa *et al.* [78] inhibition of different plant extracts against microorganisms may vary depending on plant's chemical constituents and volatility of such chemicals. Environmental and climatic variations have been cited to influence the concentrations of plant's phytochemicals thus affecting plant's antibacterial activity [77]. Antimicrobial constituents in plant extracts include compounds such as alkaloids, flavonoids, terpenoid and phenolics [79]. These antimicrobial contents have the potential to disrupt microbial cells limiting cell functions by inducing cell death

or inhibit biosynthesis of important enzymes and amino acids [80] [79]. Additionally, bioactivity of extracts from the plants could further be attributed to the hydrophobicity attributes enabling extracts to react with cell membrane and mitochondria and altering cells permeability [81] [82]. Variation in colony forming units observed is an indication that different bacteria species differ in their susceptibility to different plant extracts.

#### 4.3 Effect of Varied Concentrations of Crude Plant Extracts on Bacteria Growth

Reduction of bacteria colony forming units in all test bacteria evaluated reduced progressively with increasingly higher concentration of extracts. Minimum inhibition concentration in this study was at 0.5 ppm for all the micro-organisms studied and for all the plants extracts used. The minimum inhibition concentration of *Vangueria infausta* stem bark and crude extracts of *Physalis peruviana* leaves may be higher than 1000ppm and may be the reason why there was no inhibition in all dilution of the tested gastroenteritis pathogens. In the case of *Aspilia pluriseta*, according to reports, methanolic extracts exhibit broad-spectrum bio-activity against microbes (*Escherichia coli*, *Staphylococcus aureus*) at the concentration of 6.26 – 25 µg/mL [50]. Musa *et al.* [83] indicated a minimal inhibitory concentration of the extracts from *F. sycomorus* bark ethanolic extracts on *Salmonella typhimurium* and *E. coli* range of 6.25-1.56 mg/ml which were higher than the observed minimal inhibition concentration in this study. The concentration in this study remains lower than 50mg/ml and 300mg/ml documented earlier by Odunbaku *et al.* [76]. Variation in phytochemical activity may have been attributed to variability of environment and climate which have effect on plant phytochemical constituents. According to Li *et al.* [84] phytochemical constituents of same plant species growing under different environmental conditions can be different.

The methanolic extracts from *Erythrina abyssinnica*, *Aspilia pluriseta*, *Ficus sycomorus* performed better against *Escherichia coli*. Better performance of methanoic extract against bacteria species observed in this study are supported by other studies [85] [75] [86]. Manimozhi *et al.* [85] reported that MeOH extract of *Ficus species* exhibited better activity against all bacteria tested that included *E. coli*, *B.*

*subtilis*, and *S. aureus*. Variability of the extracts obtained by different solvent may be attributed to their differences in polarity status. Accordingly, it has been observed that Polarity of solvents has influence on the extraction efficiency of plants phytochemicals [16]. Hence, active phytochemical constituents that may not extract efficiently into acetone and hexane extract, methanol extraction is likely to provide more consistent results with wider antimicrobial activity.

### COMPETING INTERESTS

Authors have declared that no competing interests exist.

### REFERENCES

1. Degu A, Njogu P, Weru I, Karimi P. "Assessment of drug therapy problems among patients with cervical cancer at Kenyatta National Hospital, Kenya." *Gynecologic Oncology Research and Practice*. 2017;4(1):15.
2. Alhedethe A, Alhudaithy K, Zloh A. "An evaluation of prevalence of low quality of medicines in Saudi Arabia and factors associated an analytical comparative study." *Archives in Chemical Research*. 2017;2(1):1-18.
3. UN, "Transforming our world: The 2030 agenda for sustainable development," New York: United Nations, Department of Economic and Social Affairs; 2015.
4. Khuluza F, Kigera S, Heide L. "Low prevalence of substandard and falsified antimalarial and antibiotic medicines in public and faith-based health facilities of Southern Malawi." *The American Journal of Tropical Medicine and Hygiene*. 2017;96(5):1125 - 1135.
5. WHO, "Prioritization of pathogens to guide discovery, research and development of new antibiotics for drug-resistant bacterial infections, including tuberculosis (No. WHO/EMP/IAU/2017.12)," World Health Organization; 2017.
6. Nunes C, Arantes M, Pereira S, Cruz L, Passos M, Moraes L. "Plants as sources of anti-inflammatory agents." *Molecules*. 2020;25(16):3726.
7. Mir MA, Bashir N, Alfaify A, Oteef MD. "GC-MS analysis of myrtus communis extract and its antibacterial activity against gram-positive bacteria." *BioMed Central. Complementary Medicine and Therapies*. 2020;20(1):1-9.
8. Uwineza PA, Waśkiewicz A. "Recent advances in supercritical fluid extraction of natural bioactive compounds from natural plant materials." *Molecules*. 2020; 25(17):3847.
9. Sampedro J, Valdivia ER. "New antimicrobial agents of plant origin." In *Antimicrobial Compounds: Current Strategies and New Alternatives*, Villa TG and Veiga-Crespo P, Eds., Heidelberg, Germany, Springer-Verlag. 2014:83-114.
10. Altemimi A, Lakhssassi N, Baharlouei A, Watson DG, Lightfoot DA. "Phytochemicals: Extraction, isolation, and identification of bioactive compounds from plant extracts." *Plants*. 2017;6(4):42.
11. Mbuni YM, Wang S, Mwangi BN, Mbari NJ, Musili PM, Walter NO, Wang Q. "Medicinal plants and their traditional uses in local communities around Cherangani Hills, Western Kenya." *Plants*. 2020;9(3):331.
12. Cheruiyot KR, Olila D, Kateregga J. "In-vitro antibacterial activity of selected medicinal plants from Longisa region of Bomet district, Kenya." *African Health Sciences*. 2009;9(2):42-46.
13. Muthee JK, Gakuya DW, Mbaria JM, Kareru PG, Mulei CM, Njonge FK. "Ethnobotanical study of anthelmintic and other medicinal plants traditionally used in Loitokitok district of Kenya." *Journal of Ethnopharmacology*. 2011;135(1):15.
14. Gakuya DW, Itonga SM, Mbaria JM, Muthee JK, Musau JK. "Ethnobotanical survey of biopesticides and other medicinal plants traditionally used in Meru central district of Kenya." *Journal of Ethnopharmacology*. 2013;145(2):547-553.
15. Gakuya DW, Okumu MO, Kiama SG, Mbaria JM, Gathumbi PK, Mathiu PM, Nguta JM. "Traditional medicine in Kenya: Past and current status, challenges, and the way forward." *Scientific African*. 2020;8:360-367.
16. Wakeel A, Jan SA, Ullah I, Shinwari ZK, Ming X. "Solvent polarity mediates phytochemical yield and antioxidant capacity of *Isatis tinctoria*." *PeerJournal*. 2019;7:7857-7879.
17. Rafińska K, Pomastowski P, Rudnicka J, Krakowska A, Maruśka A, Narkute M, Buszewski B. "Effect of solvent and extraction technique on composition and biological activity of *Lepidium sativum*

- extracts." Food Chemistry. 2019;289:16–25.
18. Zhang H, Birch J, Pei J, Mohamed I, Ahmed A, Yang H, Dias G, Bekhit AED. "Identification of six phytochemical compounds from *Asparagus officinalis* L. root cultivars from New Zealand and China using UAE-SPE-UPLC-MS/MS: Effects of extracts on H<sub>2</sub>O<sub>2</sub>-induced oxidative stress." Nutrients. 2019;11(1):107.
  19. Sikder L, Khan M, Smrity SZ, Islam MT, Khan SA. "Phytochemical and pharmacological investigation of the ethanol extract of *Byttneria pilosa* roxb." Clinical Phytoscience. 2022;8(1):1-8.
  20. Hikmawanti NPE, Fatmawati S, Asri AW. "The effect of ethanol concentrations as the extraction solvent on antioxidant activity of katuk (*Sauropus androgynus* (L.) Merr.) leaves extracts." In IOP Conference Series: Earth and Environmental Science; 2021.
  21. Sankeshwari R, Ankola A, Bhat K, Hullatti K. "Soxhlet versus cold maceration: Which method gives better antimicrobial activity to licorice extract against *Streptococcus mutans*?" Journal of the Scientific Society. 2018;45(2):67.
  22. Abarca-Vargas R, Malacara CFP, Petricevich VL. "Characterization of chemical compounds with antioxidant and cytotoxic activities in bougainvillea x buttiana holttum and standl, (var. rose) extracts." Antioxidants. 2016;5(45):1–11.
  23. Pandey B, Thapa R, Upreti A. "Chemical composition, antioxidant and antibacterial activities of essential oil and methanol extract of *Artemisia vulgaris* and *Gaultheria fragrantissima* collected from Nepal." Asian Pacific Journal of Tropical Medicine. 2017;10(10):952–9.
  24. Shah MD, Seelan JSS, Iqbal M. "Phytochemical investigation and antioxidant activities of methanol extract, methanol fractions and essential oil of *Dillenia suffruticosa* leaves." Arabian Journal of Chemistry. 2020;13(9):7170-7182.
  25. Isah Y, Ibrahim MP. "Analysis of phytocomponents in the n-hexane extract from the stem bark of *Parinari polyandra* (benth) using FTIR and GC-MS." Bayero Journal of Pure and Applied Sciences. 2021;14(2):15-19.
  26. K. Kenya National Bureau of Statistics, "Kenya Population and Housing Census: Population by County and Sub-County," Government of Kenya, Nairobi; 2019.
  27. Kiteme MP, Onyango OB, Njagi CE, Ogolla OF. "Medicinal plants used for treatment of gastrointestinal infections in Tharaka-Nithi County." International Journal of Pathogen Research. 2022;10(4):1-11.
  28. Ejikeme C, Ezeonu CS, Eboatu AN. "Determination of physical and phytochemical constituents of some tropical timbers indigenous to Niger Delta area of Nigeria." European Scientific Journal. 2014;10(18):247-270.
  29. Hossain M, AL-Raqmi K, Al-Mijizy Z, Weli A, Al-Riyami Q. "Study of total phenol, flavonoids contents and phytochemical screening of various leaves crude extracts of locally grown *Thymus vulgaris*." Asian Pacific Journal of Tropical Biomedicine. 2013;3(9):705-710.
  30. Yadav RNS, Agarwala M. "Phytochemical analysis of some medicinal plants." Journal of Phytology. 2011;3(12):10-14.
  31. Ghosh A. De S, Dey YN. "Phytochemical investigation and chromatographic evaluation of the different extracts of tuber of *Amorphaphallus paeoniifolius* (araceae)." International Journal on Pharmaceutical and Biomedical Research. 2010;1(5):150-1.
  32. Kumar PP, Kumaravel S, Lalitha C. "Screening of antioxidant activity, total phenolics and GC-MS study of *Vitex negundo*." African Journal of Biochemistry Research. 2010;4(7):191-195.
  33. Konkolewicz D, Magenau AJ, Averick SE, Simakova A, He H, Matyjaszewski K. "ICAR ATRP with ppm Cu catalyst in water." Macromolecules. 2012;45(11):4461-4468.
  34. Lalitha M. Manual on antimicrobial susceptibility testing, Washington DC, USA: American Society for Microbiology; 2009.
  35. Puttaswamy S, Gupta SK, Regunath H, Smith LP, Sengupta S. "A comprehensive review of the present and future Antibiotic Susceptibility Testing (AST) systems." Archives of Clinical Microbiology. 2018;9(3):83-92.
  36. Andrews JM. "Determination of minimum inhibitory concentrations." Journal of Antimicrobial Chemotherapy. 2001;48(1):5-16.
  37. I. Wiegand, K. Hilpert and R. E. Hancock, "Agar and broth dilution methods to

- determine the minimal inhibitory concentration (MIC) of antimicrobial substances," *Nature Protocols*, vol. 3, no. 2, pp. 163-175, 2008.
38. R. Mogana, A. Adhikari, M. N. Tzar, R. Ramliza and C. Wiart, "Antibacterial activities of the extracts, fractions and isolated compounds from *Canarium patentinervium* Miq. against bacterial clinical isolates. *BMC Complementary Medicine and Therap*," *BMC Complementary Medicine and Therapies*, vol. 20, no. 1, pp. 1-11, 2020.
  39. P. Kuppusamy, M. M. Yusoff, N. R. Parine and N. Govindan, "Evaluation of in-vitro antioxidant and antibacterial properties of *Commelina nudiflora* L. extracts prepared by different polar solvents," *Saudi journal of biological sciences*, vol. 22, no. 3, pp. 293-301, 2015.
  40. N. Turkmen, F. Sari and Y. S. Velioglu, "Effects of extraction solvents on concentration and antioxidant activity of black and black mate tea polyphenols determined by ferrous tartrate and Folin-Ciocalteu methods," *Food chemistry*, vol. 99, no. 4, pp. 835-841, 2006.
  41. T. V. Ngo, C. J. Scarlett, M. C. Bowyer, P. D. Ngo and Q. V. Vuong, "Impact of different extraction solvents on bioactive compounds and antioxidant capacity from the root of *Salacia chinensis* L," *Journal of Food Quality*, pp. 1-8, 2017.
  42. Truong DH, Nguyen DH, Ta NTA, Bui AV, Do TH, Hoang CN. "Evaluation of the Use of Different Solvents for Phytochemical Constituents, Antioxidants, and In Vitro Anti-Inflammatory Activities of *Severinia buxifolia*. *Journal of Food Quality*. 2019;2019.
  43. Liu X, Dong M, Chen X, Jiang M, Lv X, Yan G. Antioxidant activity and phenolics of an endophytic *Xylaria* sp. from *Ginkgo biloba*. *Food chemistry*. 2007;105(2):548-54..
  44. Kadir MF, Sayeed MS, Mia MM. Ethnopharmacological survey of medicinal plants used by indigenous and tribal people in Rangamati, Bangladesh. *Journal of ethnopharmacology*. 2012 Dec 18;144(3):627-37..
  45. Mishra A, Kumar S, Bhargava A, Sharma B, Pandey AK. Studies on in vitro antioxidant and antistaphylococcal activities of some important medicinal plants. *Cellular and Molecular Biology*. 2011 Feb 12;57(1):16-25.
  46. Mishra A, Kumar S, Pandey AK. "Scientific validation of the medicinal efficacy of *Tinospora cordifolia*." *The Scientific World Journal*. 2013:1-8.
  47. Nawaz H, Waheed R, Nawaz M. "Phytochemical composition, antioxidant potential, and medicinal significance of ficus." In *Modern Fruit Industry*, I. Kahramanoglu, A. Küden and S. Çömlekçioğlu, Eds. 2019:553-573.
  48. Madivoli ES, Murigi MK, Patrick PG, Karanja GJ, Cheruiyot K, Rechab SO, Maina EG. "Phytochemical and antimicrobial evaluation of selected medicinal plants in Meru community of Kenya. *Journal of Medicinal Plants for Economic Development*. 2018;2(1):1-4.
  49. Osama A, Awdelkarim S. "Phytochemical screening of *Ficus sycomorus* L. bark and *Cleome gynandra* L. aerial parts." *Journal of Pharmacognosy and Phytochemistry*. 2015;4(4):24-27.
  50. Njerua SN, Muema JM. "Antimicrobial activity, phytochemical characterization and gas chromatography-mass spectrometry analysis of *Aspilia pluriseta* Schweinf. Extracts." *Heliyon*. 2020; 6(10):5195-5165.
  51. Musyimi DM, Ogur JA, Muema PM. "Effects of leaf and root extracts of aspilia plant (*Aspilia mossambicensis*) (Oliv) Wild. on some selected microorganisms," *International Journal of Biological Chemistry*. 2007;1(4):213-220.
  52. Nzuki DM. "Utilization of herbal medicine among children under 5 years of age in Tharaka Nithi County, Kenya." *Doctoral Dissertation, Kenyatta University, Kenya*; 2016.
  53. Obakiro SB, Kiprop A, Kigundu E, K'Owino I, Odero MP, Manyim S, Bunalema L. "Traditional medicinal Uses, phytoconstituents, bioactivities, and toxicities of *Erythrina abyssinica* Lam. ex DC. (fabaceae): A systematic review." *Evidence-Based Complementary and Alternative Medicine*. 2021:1-43.
  54. Manyarara TE, Chifamba J, Tarugarira FT. "Antifungal activity of *Ziziphus mucronata* and *Erythrina abyssinica* bark crude extracts on *Cryptococcus neoformans* and *Candida albicans* species." *Journal of Pharmaceutical Research International*. 2016;10(3):1-11.
  55. Desai SD, Desai DG, Kaur H. "Saponins and their biological activities." *Pharma Times*. 2009;41(3):13-16.

56. Heim KE, Tagliaferro AR, Bobilya DJ. "Flavonoid antioxidants: Chemistry, metabolism and structure-activity relationships." *The Journal of Nutritional Biochemistry*. 2002;13(10):572-584.
57. Pandey AK, Mishra AK, Mishra A. "Antifungal and antioxidative potential of oil and extracts derived from leaves of Indian spice plant *Cinnamomum tamala*." *Cellular and Molecular Biology*. 2012;58(1):142-147.
58. Meng XH, Liu C, Fan R, Zhu LF, Yang SX, Zhu HT, Zhang YJ. "Antioxidative flavan-3-ol dimers from the leaves of *Camellia fangchengensis*." *Journal of Agricultural and Food Chemistry*. vol. 2018;66(1):247-254.
59. Kumar S, Pandey AK. "Chemistry and biological activities of flavonoids: An overview." *The Scientific World Journal*. 2013:1-16.
60. Cazarolli LH, Zanatta L, Alberton EH, Bonorino Figueiredo MSR, Folador P, Damazio RG, Barreto Silva FRM. "Flavonoids: Prospective drug candidates. Mini reviews in medicinal chemistry." 2008;8(13):1429-1440.
61. Singh R, Singh S, Kumar S, Arora S. "Evaluation of antioxidant potential of ethyl acetate extract/fractions of *Acacia auriculiformis*." *Food and Chemical Toxicology*. 2007;45(7):1216-1223.
62. Han X, Shen T, Lou H. "Dietary polyphenols and their biological significance." *International Journal of Molecular Sciences*. 2007;8(9):950-988.
63. Ali SS, Kasoju N, Luthra A, Singh A, Sharanabasava H, Sahu A, Bora U. "Indian medicinal herbs as sources of antioxidants." *Food Research International*. 2008;41(1):1-15.
64. Canini A, Alesiani D, D'Arcangelo G, Tagliatesta P. "Gas chromatography–mass spectrometry analysis of phenolic compounds from *Carica papaya* L. leaf." *Journal of Food Composition and Analysis*. 2007;20(7):584-590.
65. Epand RFSPB, Epand RM. "Bacterial lipid composition and the antimicrobial efficacy of cationic steroid compounds (ceragenins)." *Biochimica et Biophysica Acta (BBA)-Biomembranes*. 2007;1768(10):2500-2509.
66. Kukula-Koch WA, Widelski J. "Alkaloids," in pharmacognosy, Badal S and Delgoda R, Eds., London, UK., Academic Press. 2017:163-198.
67. Ayoola GA, Coker HA, Adesegun SA, Adepoju-Bello AA, Obaweya K, Ezennia EC, Atangbayila TO. "Phytochemical screening and antioxidant activities of some selected medicinal plants used for malaria therapy in Southwestern Nigeria." *Tropical Journal of Pharmaceutical Research*. 2008;7(3):1019-1024.
68. Ahmad I, Mehmood Z, Mohammad F. "Screening of some Indian medicinal plants for their antimicrobial properties." *Journal of Ethnopharmacology*. 1998;62:183-9.
69. Lu LWS, Liu S, Jiang B, Wu SG. "Tannin inhibits HIV-1 entry by targeting gp41." *Acta Pharmacologica Sinica*. 2004;25(2):213-218.
70. Funatogawa K, Hayashi S, Shimomura H, Yoshida T, Hatano T, Ito HHY. "Antibacterial activity of hydrolyzable tannins derived from medicinal plants against *Helicobacter pylori*." *Microbiology and Immunology*. 2004;48(4):251-261.
71. Awuchi CG. "The biochemistry, toxicology, and Uses of the acologically active phytochemicals: Alkaloids, terpenes, polyphenols, and glycosides." *Merit Research Journals*. 2020;5(1):6-21.
72. Njoku VO, Obi C. "Phytochemical constituents of some selected medicinal plants." *African Journal of Pure and Applied Chemistry*. 2009;3(11):228-233.
73. Korir K, Bii C, Kiiyukia C, Mutai C. "Antimicrobial activities of *Clutia abyssinica* and *Erythrina abyssinica* plants extracts used among the Kipsigis community of Bomet district in Kenya." *Natural Products*. 2011;7(5):247-252.
74. Njeru SN, Obonyo MA. "Potency of extracts of selected plant species from Mbeere, Embu County-Kenya against *Mycobacterium tuberculosis*." *Journal of Medicinal Plants Research*. vol. 2016;10(12):149-157.
75. Salem MZ, Salem AZ, Camacho LM, Ali HM. "Antimicrobial activities and phytochemical composition." *African Journal of Microbiology Research*. 2013;7(33):4207-4219.
76. Odunbaku O, Ilusanya O, Akasoro K. "Antibacterial activity of ethanolic leaf extract of *Ficus exasperata* on *Escherichia coli* and *Staphylococcus albus*." *Scientific Research and Essay*. 2008;3(11):562-564.
77. Chilufya G, Masaiti G, Malambo C, Mudenda S. "Antibacterial properties of *Ficus sycomorus* bark extract against *Staphylococcus aureus* and *Escherichia*

- coli*." International Journal of Biomedical Investigation. 2019;2(121):1-8.
78. Mostafa AA, Al-Askara AA, Almaary KS, Dawoud TM, Sholkamy EN, Bakric MM. "Antimicrobial activity of some plant extracts against bacterial strains causing food poisoning diseases." Saudi Journal of Biological Sciences. 2018;25(2):361-366.
79. Alibi S, Dámaso C, Navas J. "Plant-derivatives small molecules with antibacterial activity." Antibiotics. 2021; 10(3):23-30.
80. Khameneh B, Iranshahy M, Soheili V, Bazzaz BS. "Review on plant antimicrobials: A mechanistic viewpoint." Antimicrobial Resistance and Infection Control. 2019;8(1):1-28.
81. Friedman M, Henika P, Levin C, Mandrell R. "Antibacterial activities of plant essential oils and their components against *Escherichia coli* O157:H7 and *Salmonella enterica* in apple juice." Journal of Agricultural and Food Chemistry. 2004;52:6042-6048.
82. Tiwari B, Valdramidi V, O'Donnell C, Muthukumarappan K, Bourke P, Cullen P. "Application of natural antimicrobials for food preservation." Journal of Agricultural and Food Chemistry. 2009;57:5987-6000.
83. Musa I, Manga SS, Muhammad A, Tondi ZY, Saadu M. "The *In vitro* antibacterial effects of ficus sycomorus stem bark extracts on *Salmonella typhimurium* and *Escherichia coli*." BiblioMed. 2020;7(1): 108-113.
84. Li Y, Kong D, Fu Y, Sussman MR, Wu H. "The effect of developmental and environmental factors on secondary metabolites in medicinal plants," Plant Physiology and Biochemistry, vol. 148, pp. 80-89, 2020.
85. Manimozhi D, Sankaranarayanan S, Sampathkumar G, "Evaluating the antibacterial activity of flavonoids extracted from ficus benghalensis." International Journal of Pharmaceutical and Biological Research. 2012;3(1).
86. Mudzengi C, Murwira A, Tivapasic M, Murungweni C, Burumue J, Halimani T. "Antibacterial activity of aqueous and methanol extracts of selected species used in livestock health management." Pharmaceutical Biology. 2017;55:1054-1060.

© 2023 Kiteme et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:  
<https://www.sdiarticle5.com/review-history/98790>