

CHUKA



UNIVERSITY

UNIVERSITY EXAMINATIONS

EXAMINATION FOR THE AWARD OF DEGREE OF BACHELOR OF SCIENCE IN  
PHYSICS AND BACHELOR OF EDUCATION SCIENCE

PHYS 414: NUCLEAR AND ATOMIC PHYSICS

STREAMS: BSC (PHYS), BED (SCI)

TIME: 2 HOURS

DAY/DATE: TUESDAY 04/12/2018

11.30 A.M. – 1.30 P.M.

INSTRUCTIONS:

- Answer question ONE and any other TWO questions.

QUESTION ONE (30 MARKS)

- (a) Define the terms nuclear fusion and nuclear fission. (2 marks)
- (b) The electron in the hydrogen atom makes transition from the  $n = 4$  to the ground state. Find the wavelength and frequency of the emitted photon. ( $R_H = 1.097 \times 10^7 M^{-1}$ ). (4 marks)
- (c) Show that  $E_n = \frac{-m_e k_e^2 e^4}{2h^2} \left(\frac{1}{n^2}\right) n = 1, 2, 3 \dots$  (4 marks)
- (d) State Pauli's Exclusion principle. (2 marks)
- (e) Write electronic configuration of  $z = 21$  utilizing exclusion principle. (3 marks)
- (f) Estimate the energy of the characteristic X-ray emitted from a tungsten target when an electron drops from a N shell ( $n=4$ ) to a vacancy in the K shell ( $n = 1$ ).  $Z = 74$ . (5 marks)
- (g) Draw an energy level diagram for hydrogen and at least show four series. (4 marks)
- (h) The nuclear reaction  ${}_0^1n + {}_5^{10}B \rightarrow {}_3^7Li + {}_2^4He$  is observed to occur even when very slow moving neutrons ( $M_n = 1.0087u$ ) strike a boron atom at rest. For a particular reaction in which  $KE=0$ , the helium ( $M_{He}=4.0026u$ ) is observed to have a speed of  $9.30 \times 10^6$  m/s.

Determine

- (i) The KE of the lithium ( $M_{Li} = 7.0160$ ) (4 marks)
- (ii) The Q value of the reaction (2 marks)

**QUESTION TWO (20 MARKS)**

- (a) A “clever” technician decides to heat some water for his coffee with an x-ray machine. If the machine produces 10 rad/s, how long will it take to raise the temperature of a cup of water by  $50^{\circ}C$ ? Ignore heat losses during this time? (6 marks)
- (b) In a certain experiment,  $0.024 \mu Ci$  of  ${}_{15}^{32}P$  is injected into a medium containing a culture of bacteria. After 2 hours the cells are washed and a detector that is 70% efficient (counts 70% of emitted rays) records 1440 counts per minute from all the cells. What percentage of the original was taken up by the cells? (3 marks)
- (c) Show that half-life of a radioactive material can be expressed as  $t_{\frac{1}{2}} = \frac{0.693}{\lambda}$  where  $\lambda$  = decay constant (3 marks)
- (iii) Why is the quantity of energy different in the two (1 mark)
- (d) An isotope of an element radon has half-life of 8 days, a sample of radon originally contains  $8.2 \times 10^{16}$  atoms, take one day to be  $86 \times 10^3$  seconds, calculate
  - (i) The number of radon atoms remaining after 32 days. (2 marks)
  - (ii) The rate of decay of the radon sample after 32 days. (2 marks)
- (e) Define the following
  - (i) LASER
  - (ii) MASER
  - (iii) Phosphorescence (3 marks)

**QUESTION THREE (20 MARKS)**

- (a) State any four useful applications of radioactivity (4 marks)
- (b) Calculate the binding energy in alpha particle (Helium-4) nucleus in MeV. Take  
 Mass of neutron = 1.008665u  
 Mass of helium nucleus = 4.001508u  
 Mass of a photon = 1.007276u (3 marks)
- (c) State and explain briefly FIVE types of stationary power reactors (5 marks)
- (d) State Neil Bohr’s atomic model postulates (3 marks)
- (e) State five Hazard of radioactivity (5 marks)

**QUESTION FOUR**

- (a) Natural gold has only one isotope,  $^{197}_{79}\text{Au}$ . If gold is bombard with slow neutron,  $\bar{\nu}$  particles are emitted.
- Write the appropriate reaction equation.
  - Calculate the maximum energy of the emitted beta particles. The mass of  $^{198}_{80}\text{Hg}$  is 197.966 75 u. (4 marks)
- (b) Sketch a graph showing the average binding energy per nucleon as a function of mass number A. (4 marks)
- (c) Estimate the temperature required for a deuterium-tritium fusion (d-t) to occur  $R_d=1.5$  fm and  $r_t=1.7$  fm. (4 marks)
- (d) An animal bone fragment found in archaeological site has a carbon mass of 400g. It registers an activity of 20 decays/s. What is the age of the bone? (Ratio of C-14:C-12 when the animal was alive was  $1.3 \times 10^{-12}$ ) (4 marks)
- (e) Draw a well labeled diagram of a nuclear reactor. (4 marks)

**QUESTION FIVE**

- (a) Calculate the energy in MeV liberated when helium is produced.
- by fusing two neutrons and two protons (3 marks)  
 $m_p = 1.007825u, M_n = 1.008665u$
  - by fusing two deuterium nuclei  $^2_1\text{H} = 2.014102$  (3 marks)
  - Why the difference? (2 marks)
- (b) Calculate the total binding energy and the average binding energy per nucleon for  $^{56}_{26}\text{Fe}$  the most common stable isotope of Iron ( $p=1.007825u, n=1.008665u$  and  $M_{\text{Fe}}=55.9349u$ ) (5 marks)
- (c) Compare at least THREE properties of alpha, beta and gamma decays. (3 marks)
- (d) Describe the kind of decay particle in the following nuclear equations A, B, C and D
- $^1_0n + ^{238}_{92}\text{U} \rightarrow ^{239}_{92}\text{U} + A$
  - $^2_1\text{H} + ^{14}_7\text{N} \rightarrow ^{12}_6\text{C} + B$
  - $^{212}_{83}\text{Bi} \rightarrow ^{208}_{81}\text{Tl} + C$
  - $^2_1\text{H} + ^2_1\text{H} \rightarrow D$  (4 marks)
-